

PATIENT DEMOGRAPHICS

Patient Name: DOB: Phone: Address: City/ST/Zip: Allergies: Patient Status: [] New to Therapy [] Dose or Frequency Change [] Order Renewal Sex: [] Male [] Female [] Unknown

INSURANCE INFORMATION: Please attach copy of insurance card (front and back).

DIAGNOSIS*

*ICD 10 Code Required

[] Nausea, ICD10 R11.0 [] Vomiting, unspecified, ICD10 R11.10 [] Nausea and vomiting, unspecified, ICD10 R11.2 [] Encounter for antineoplastic chemotherapy, ICD10 Z51.11 [] Adverse effect of antineoplastic and immunosuppressive drugs, ICD10 T45.1X5A [] Adverse effect of antineoplastic and immunosuppressive drugs, ICD10 T45.1X5D [] Adverse effect of antineoplastic and immunosuppressive drugs, ICD10 T45.1X5S [] Other: ICD10:

PRESCRIPTION INFORMATION

Table with 4 columns: MEDICATION, DOSE, DIRECTIONS/DURATION, QTY. Row 1: Sancuso® (Granisetron Transdermal System), 1 transdermal system delivering granisetron 3.1 mg / 24 hours, Apply a single transdermal system to the upper outer arm 24-48 hours before chemotherapy. A single transdermal system can be worn for up to 7 days. # of Patches: # of Refills:

Is patient currently receiving therapy above from another facility?

[] NO [] YES

If yes, Facility Name:

Date of last treatment: Date of next treatment:

PRESCRIBER INFORMATION

Prescriber's Signature: Date: Prescriber's Name: Provider NPI: Specialty: Address: City/ST/Zip: Contact Person: Phone #: Fax #: Email Where Follow Up Documentation Should Be Sent:

REQUIRED CLINICAL DOCUMENTATION

Please attach medical records: Initial H&P, current MD progress notes, medication list, and labs/test results to support diagnosis.

Clinical Information, select all that apply:

[] Sancuso® (granisetron) will be administered for: [] Prevention of chemotherapy-induced nausea and vomiting [] Treatment of breakthrough nausea and/or vomiting due to chemotherapy [] The patient will be receiving moderately and/or highly emetogenic chemotherapy for up to five consecutive days. Please specify chemotherapy regimen: [] The patient is at risk for chemotherapy-induced nausea or vomiting. Please specify: [] Young age (<55 years) [] Female sex [] Previous history of chemotherapy induced nausea or vomiting [] Little or no previous alcohol use [] History of motion sickness or morning sickness during pregnancy [] High anxiety [] Other: [] The patient is unable to swallow or digest tablets/capsules.

PRIOR FAILED THERAPIES FOR NAUSEA/VOMITING

Medication Failed: Dates of Treatment: Reason for D/C: Medication Failed: Dates of Treatment: Reason for D/C: Medication Failed: Dates of Treatment: Reason for D/C: Medication Failed: Dates of Treatment: Reason for D/C: